

## **APPENDIX B**

# **SIGNIFICANT COASTAL FISH AND WILDLIFE HABITATS**

COASTAL FISH & WILDLIFE HABITAT RATING FORM

---

Name of Area: **Haverstraw Bay**  
 Designated: **November 15, 1987**  
 County(ies): **Rockland; Westchester**  
 Town(s): **Clarkstown, Haverstraw, Stony Point; Cortlandt**  
 7½' Quadrangle(s): **Haverstraw, NY; NOAA Chart No. 12343**

---

Score    Criterion

- 40      Ecosystem Rarity (ER)  
           The most extensive area of shallow estuarine habitat in the lower Hudson River (and in New York State), but rarity reduced by human disturbances; geometric mean:  $(25 \times 64)^{1/2} = 40$ .
- 36      Species Vulnerability (SV)  
           Shortnose sturgeon (E) regularly occur in the area.
- 38      Human Use (HU)  
           The area contributes to recreational and commercial fisheries throughout the northeastern U.S.;  
           additive division:  $25 + 25/2 = 38$
- 25      Population Level (PL)  
           A major spawning, nursery, and wintering area for various estuarine fish species; population levels unusual in the northeastern U.S.
- 1.2     Replaceability (R)  
           Irreplaceable.

---

SIGNIFICANCE VALUE =  $[(ER + SV + HU + PL) \times R] = 166$



## **SIGNIFICANT COASTAL FISH AND WILDLIFE HABITATS PROGRAM A PART OF THE NEW YORK COASTAL MANAGEMENT PROGRAM**

### **BACKGROUND**

New York State's Coastal Management Program (CMP) includes a total of 44 policies which are applicable to development and use proposals within or affecting the State's coastal area. Any activity that is subject to review under Federal or State laws, or under applicable local laws contained in an approved local waterfront revitalization program will be judged for its consistency with these policies.

Once a determination is made that the proposed action is subject to consistency review, a specific policy aimed at the protection of fish and wildlife resources of statewide significance applies. The specific policy statement is as follows: "Significant coastal fish and wildlife habitats will be protected, preserved, and, where practical, restored so as to maintain their viability as habitats." The New York State Department of Environmental Conservation (DEC) evaluates the significance of coastal fish and wildlife habitats, and following a recommendation from the DEC, the Department of State designates and maps specific areas. Although designated habitat areas are delineated on the coastal area map, the applicability of this policy does not depend on the specific location of the habitat, but on the determination that the proposed action is subject to consistency review.

Significant coastal fish and wildlife habitats are evaluated, designated and mapped under the authority of the Coastal Management Program's enabling legislation, the Waterfront Revitalization and Coastal Resources Act (Executive Law of New York, Article 42). These designations are subsequently incorporated in the Coastal Management Program under authority provided by the Federal Coastal Zone Management Act.

This narrative constitutes a record of the basis for this significant coastal fish and wildlife habitat's designation and provides specific information regarding the fish and wildlife resources that depend on this area. General information is also provided to assist in evaluating impacts of proposed activities on parameters which are essential to the habitat's values. This information is to be used in conjunction with the habitat impairment test found in the impact assessment section to determine whether the proposed activities are consistent with this policy.

## **DESIGNATED HABITAT: HAVERSTRAW BAY**

### **HABITAT DESCRIPTION:**

Haverstraw Bay extends approximately six miles on the Hudson River, from Stony Point to Croton Point, in the Towns of Stony Point, Haverstraw, and Clarkstown, in Rockland County, and the Town of Cortlandt, in Westchester County (7.5' Quadrangle: Haverstraw, N.Y.; NOAA Chart No. 12343).

The fish and wildlife habitat encompasses the entire river over this approximate six mile reach, which is the widest section of the Hudson estuary. Haverstraw Bay has extensive shallow areas (less than 15 feet deep at mean low water) which deepen to a navigation channel (which is dredged to maintain a depth of about 35 feet) in the western half of the area. During much of the year, this area is the place where freshwater from the upper river mixes with salt water from the Atlantic, producing a predominantly brackish water habitats, with salinities varying from 0-10 ppt. The land area surrounding Haverstraw Bay supports a variety of land uses, including industrial, commercial, residential, and recreational developments, although much undeveloped forestland also remains. Habitat disturbances, such as dredging, shoreline filling and bulkheading, waste disposal, and pollution from upland and in-river sources, have all been significant at some time during the recent history of this area.

### **FISH AND WILDLIFE VALUES:**

Despite various habitat disturbances, Haverstraw Bay possesses a combination of physical and biological characteristics that make it one of the most important fish and wildlife habitats in the Hudson River estuary. The regular occurrence of brackish water over extensive areas of shallow bottom creates highly favorable (if not essential) conditions for biological productivity within the estuary, including submergent vegetation, phytoplankton and zooplankton, aquatic invertebrates, and many fish species.

Although the location of the salt front varies annually (and seasonally), Haverstraw Bay regularly comprises a substantial part of the nursery area for striped bass, American shad, white perch, tomcod, and Atlantic sturgeon that are produced in the Hudson. Other anadromous species, such as blueback herring and alewife, spawn in upstream freshwater areas, but move south and concentrate in this area before leaving the river in the fall.

Haverstraw Bay is also a major nursery and feeding area for certain marine species, most notably bay anchovy, Atlantic menhaden, and blue claw crab. Depending on location of the salt front, a majority of the spawning and wintering populations of Atlantic sturgeon in the Hudson may reside in Haverstraw Bay. Shortnose sturgeon (E) usually winter in this area as well. Significant numbers of waterfowl may occur in Haverstraw Bay during spring (March-April) and fall (September-November) migrations, but the extent of this use is not well documented.

Haverstraw Bay is a critical habitat for most estuarine-dependent fisheries originating from the Hudson River. This area contributes directly to the production of in-river and ocean populations of food, game, and forage fish species. Consequently, commercial and recreational fisheries throughout the North Atlantic depend on, or benefit from, these biological inputs from the Hudson River estuary.

#### IMPACT ASSESSMENT:

A **habitat impairment test** must be met for any activity that is subject to consistency review under federal and State laws, or under applicable local laws contained in an approved local waterfront revitalization program. If the proposed action is subject to consistency review, then the habitat protection policy applies, whether the proposed action is to occur within or outside the designated area.

The specific **habitat impairment test** that must be met is as follows.

In order to protect and preserve a significant habitat, land and water uses or development shall not be undertaken if such actions would:

- destroy the habitat; or,
- significantly impair the viability of a habitat.

*Habitat destruction* is defined as the loss of fish or wildlife use through direct physical alteration, disturbance, or pollution of a designated area or through the indirect effects of these actions on a designated area. Habitat destruction may be indicated by changes in vegetation, substrate, or hydrology, or increases in runoff, erosion, sedimentation, or pollutants.

*Significant impairment* is defined as reduction in vital resources (e.g., food, shelter, living space) or change in environmental conditions (e.g., temperature, substrate, salinity) beyond the tolerance range of an organism. Indicators of a significantly impaired habitat focus on ecological alterations and may include but are not limited to reduced carrying capacity, changes in community structure (food chain relationships, species diversity), reduced productivity and/or increased incidence of disease and mortality.

The *tolerance range* of an organism is not defined as the physiological range of conditions beyond which a species will not survive at all, but as the ecological range of conditions that supports the species population or has the potential to support a restored population, where practical. Either the loss of individuals through an increase in emigration or an increase in death rate indicates that the tolerance range of an organism has been exceeded. An abrupt increase in death rate may occur as an environmental factor falls beyond a tolerance limit (a range has both upper and lower limits). Many environmental factors, however, do not have a sharply defined tolerance limit, but produce increasing emigration or death rates with increasing departure from conditions that are optimal for the species.

The range of parameters which should be considered in applying the habitat impairment test include but are not limited to the following:

1. physical parameters such as living space, circulation, flushing rates, tidal amplitude, turbidity, water temperature, depth (including loss of littoral zone), morphology, substrate type, vegetation, structure, erosion and sedimentation rates;
2. biological parameters such as community structure, food chain relationships, species diversity, predator/prey relationships, population size, mortality rates, reproductive rates, meristic features, behavioral patterns and migratory patterns; and,
3. chemical parameters such as dissolved oxygen, carbon dioxide, acidity, dissolved solids, nutrients, organics, salinity, and pollutants (heavy metals, toxics and hazardous materials).

Although not comprehensive, examples of generic activities and impacts which could destroy or significantly impair the habitat are listed below to assist in applying the habitat impairment test to a proposed activity.

Any activity that would substantially degrade water quality, increase turbidity or sedimentation, or alter water salinities or temperatures in Haverstraw Bay would result in significant impairment of the habitat. Any physical modification of the habitat or adjacent wetlands, through dredging, filling, or bulkheading, would result in a direct loss of valuable habitat area.

Habitat disturbances would be most detrimental during fish spawning and early developmental periods, which generally extend from April through August for most anadromous species using the area. Discharges of sewage or stormwater runoff containing sediments or chemical pollutants may result in significant adverse impacts on fish populations. Similarly, spills of oil or other hazardous substances, and leachate of contaminated groundwater, constitute a potential threat to fish and wildlife in the bay. Of particular concern in this major estuarine system are the potential effects of hydrologic disturbances, and effluent discharges. Existing areas of natural vegetation bordering Haverstraw Bay should be maintained to provide soil stabilization and buffer areas.

KNOWLEDGEABLE CONTACTS:


Tom Hart  
N.Y.S. Department of State  
Division of Coastal Resources &  
Waterfront Revitalization  
162 Washington Avenue  
Albany, NY 12231  
Phone: (518) 474-6000

Wayne Elliot, Fisheries Manager  
or Jack Isaacs, Environmental Protection Biologist  
NYSDEC - Region 3  
21 South Putt Corners Road  
New Paltz, NY 12561  
Phone: (914) 255-5453

Doug Carlson, Hudson River Fisheries Unit  
NYSDEC - Region 4  
Route 10, Jefferson Road  
Stamford, NY 12167  
Phone: (607) 652-7364


William L. Dovel  
Route 1, Box 800  
Venice, FL 33595  
Phone: (813) 493-4354

NYSDEC Information Services  
700 Troy-Schenectady Road  
Latham, NY 12110  
Phone: (518)783-3932



**Significant Coastal Fish and Wildlife Habitats**

Haverstraw Bay (In part)  
Part 1 of 2



**Significant Coastal Fish and Wildlife Habitats**

Croton River & Bay  
 Haverstraw Bay (In part)  
 Part 2 of 2

New York State  
 Department of State  
 Division of  
 Coastal Resources

